

LOW-TEMPERATURE HIGH-RESOLUTION INFRARED SPECTRUM OF ETHANE-1D, C₂H₅D: ROTATIONAL ANALYSIS OF THE ν_{17} BAND NEAR 805 cm⁻¹ using ERHAM.

PETER GRONER, *Department of Chemistry, University of Missouri - Kansas City, Kansas City, MO, USA*; ADAM M DALY, BRIAN DROUIN, JOHN PEARSON, KEEYOON SUNG, LINDA BROWN, *Jet Propulsion Laboratory, California Institute of Technology, Pasadena, CA, USA*; ARLAN MANTZ, *Department of Physics, Astronomy and Geophysics, Connecticut College, New London, CT, USA*; MARY ANN H. SMITH, *Science Directorate, NASA Langley Research Center, Hampton, VA, USA*.

The high-resolution infrared spectrum of gaseous ethane-d₁ at 130 K shows transitions that are split into A and E components due to the interaction of overall rotation with the internal rotation of the CH₃ group. An analysis of the spectrum from 680 to 900 cm⁻¹ with an expanded version of the program ERHAM^{a,b} is in progress to assign the bands at E(ν_{17}) = 805 cm⁻¹ and E(ν_{11}) = 715 cm⁻¹. A discussion of the interactions among the fundamental levels of ν_{17} and ν_{11} with overtone levels of ν_{18} and the(CH₃ torsion) will be given. ERHAM has been and continues to be very successful in the analysis of pure the rotational spectra of molecules containing internal rotation and the vibrational spectrum of C₂H₅D serves as an excellent system to test the extension of the program.

^aP. Groner, *J. Chem. Phys.* **107** 4483 (1997)

^bP. Groner, *J. Mol. Spectrosc.* **278** 52 (2012)